

Referenties

1. Earl AM, Losick R, Kolter R: Ecology and genomics of *Bacillus subtilis*. *Trends Microbiol* 2008, 16:269–275.
2. Harwood CR: *Bacillus subtilis* and its relatives: molecular biological and industrial workhorses. *Trends Biotechnol* 1992, 10:247–256.
3. Kunst F, Ogasawara N, Moszer I, Albertini AM, Alloni G, Azevedo V, et al: The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* 1997, 390:249–256.
4. Barbe V, Cruveiller S, Kunst F, Lenoble P, Meurice G, Sekowska A, et al: From a consortium sequence to a unified sequence: the *Bacillus subtilis* 168 reference genome a decade later. *Microbiology* 2009, 155:1758–1775.
5. Kobayashi K, Ehrlich SD, Albertini A, Amati G, Andersen KK, Arnaud M, et al: Essential *Bacillus subtilis* genes. *Proc Natl Acad Sci USA* 2003, 100:4678–4683.
6. Nicolas P, Mäder U, Dervyn E, Rochat T, Leduc A, Pigeonneau N, et al: Condition-dependent transcriptome reveals high-level regulatory architecture in *Bacillus subtilis*. *Science* 2012, 335:1103–1106.
7. Büscher JM, Liebermeister W, Jules M, Uhr M, Muntel J, Botella E, et al: Global network reorganization during dynamic adaptations of *Bacillus subtilis* metabolism. *Science* 2012, 335:1099–1103.
8. Otto A, Bernhardt J, Meyer H, Schaffer M, Herbst FA, Siebourg J, et al: Systems-wide temporal proteomic profiling in glucose-starved *Bacillus subtilis*. *Nat Commun* 2010, 1:137.
9. Becher D, Büttner K, Moche M, Hessling B, Hecker M: From the genome sequence to the protein inventory of *Bacillus subtilis*. *Proteomics* 2011, 11:2971–2980.
10. Maass S, Sievers S, Zühlke D, Kuzinski J, Sappa PK, Muntel J, et al: Efficient, global-scale quantification of absolute protein amounts by integration of targeted mass spectrometry and two-dimensional gel-based proteomics. *Anal Chem* 2011, 83:2677–2684.

11. Tjalsma H, Bolhuis A, Jongbloed JD, Bron S, van Dijl JM: Signal peptide β -dependent protein transport in *Bacillus subtilis*: a genome-based survey of the secretome. *Microbiol Mol Biol Rev* 2000, 64:515–547.
12. Antelmann H, Tjalsma H, Voigt B, Ohlmeier S, Bron S, van Dijl JM, et al: A proteomic view on genome-based signal peptide predictions. *Genome Res* 2001, 11:1484–1502.
13. Tjalsma H, Antelmann H, Jongbloed JD, Braun PG, Darmon E, Dorenbos R, et al: Proteomics of protein secretion by *Bacillus subtilis*: separating the 'secrets' of the secretome. *Microbiol Mol Biol Rev* 2004, 68:207–233.
14. Jongbloed JDH, Martin U, Antelmann H, Hecker M, Tjalsma H, Venema G, et al: TatC is a specificity determinant for protein secretion via the twin β arginine translocation pathway. *J Biol Chem* 2000, 275:41350–41357.
15. Wolff S, Antelmann H, Albrecht D, Becher D, Bernhardt J, Bron S, et al: Towards the entire proteome of the model bacterium *Bacillus subtilis* by gel-based and gel-free approaches. *J Chromatogr B Analyt Technol Biomed Life Sci* 2007, 849:129–140.
16. Antelmann H, Darmon E, Noone D, Veening JW, Westers H, Bron S, et al: The extracellular proteome of *Bacillus subtilis* under secretion stress conditions. *Mol Microbiol* 2003, 49:143–156.
17. Darmon E, Dorenbos R, Meens J, Freudl R, Antelmann H, Hecker M, et al: A disulfide bond-containing alkaline phosphatase triggers a BdbC β -dependent secretion stress response in *Bacillus subtilis*. *Appl Environ Microbiol* 2006, 72:6876–6885.
18. Tjalsma H, van Dijl JM: Proteomics-based consensus prediction of protein retention in a bacterial membrane. *Proteomics* 2005, 5:4472–4482.
19. Brockmeier U, Caspers M, Freudl R, Jockwer A, Noll T, Eggert T: Systematic screening of all signal peptides from *Bacillus subtilis*: a powerful strategy in optimizing heterologous protein secretion in Gram-positive bacteria. *J Mol Biol* 2006, 362:393–402.
20. Dalbey RE, Wang P, van Dijl JM: Membrane proteases in the bacterial protein secretion and quality control pathway. *Microbiol Mol Biol Rev* 2012, 76:311–330.

21. van Roosmalen ML, Geukens N, Jongbloed JDH, Tjalsma H, Dubois J-YF, Bron S, et al: Type I signal peptidases of Gram-positive bacteria. *Biochim Biophys Acta* 2004, 1694:279–297.
22. Saito A, Hizukuri Y, Matsuo E, Chiba S, Mori H, Nishimura O, et al: Post-translocation cleavage of signal peptides is catalyzed by the site-2 protease (S2P) in bacteria. *Proc Natl Acad Sci USA* 2011, 108:13740–13745.
23. Yuan J, Zweers JC, van Dijl JM, Dalbey RE: Protein transport across and into cell membranes in bacteria and archaea. *Cell Mol Life Sci* 2010, 67:179–199.
24. Sarvas M, Harwood CR, Bron S, van Dijl JM: Post-translocational folding of secretory proteins in Gram-positive bacteria. *Biochim Biophys Acta* 2004, 1694:311–327.
25. Jongbloed JDH, van der Ploeg R, van Dijl JM: Bifunctional TatA subunits in minimal Tat protein translocases. *Trends Microbiol* 2006, 14:2–4.
26. Fröbel J, Rose P, Lausberg F, Blümmel AS, Freudl R, Müller M: Transmembrane insertion of twin-arginine signal peptides is driven by TatC and regulated by TatB. *Nat Commun* 2012, 3:1311.
27. Rollauer SE, Tarry MJ, Graham JE, Jääskeläinen M, Jäger F, Johnson S, et al: Structure of the TatC core of the twin-arginine protein transport system. *Nature* 2012, 492:210–214.
28. van der Ploeg R, Mäder U, Homuth G, Schaffer M, Denham EL, Monteferrante CG, et al: Environmental salinity determines the specificity and need for Tat-dependent secretion of the YwbN protein in *Bacillus subtilis*. *PLoS One* 2011, 6:e18140.
29. van der Ploeg R, Barnett JP, Vasisht N, Goosens VJ, Poether DC, Robinson C, et al: Salt-sensitivity of Minimal Twin-arginine Translocases. *J Biol Chem* 2011, 286:43759–43770.
30. Monteferrante CG, MacKichan C, Marchadier E, Prejean M-V, Carballido-López R, Van Dijl JM: Mapping the twin-arginine protein translocation network of *Bacillus subtilis*. *Proteomics* 2012, in press.
31. van Dijl JM, Braun PG, Robinson C, Quax WJ, Antelmann H, Hecker M, et al: Functional genomic analysis of the *Bacillus subtilis* Tat pathway for

- protein secretion. *J Biotechnol* 2002, 98:243–254.
32. van der Ploeg R, Monteferante CG, Piersma S, Barnett JP, Kouwen TRHM, Robinson C, et al: High salinity growth conditions promote Tat independent secretion of Tat substrates in *Bacillus subtilis*. *Appl Environ Microbiol* 2012, 78:7733–7744.
33. Barnett JP, Eijlander RT, Kuipers OP, Robinson C: A minimal Tat system from a gram-positive organism: a bifunctional TatA subunit participates in discrete TatAC and TatA complexes. *J Biol Chem* 2008, 283:2534–2542.
34. Barnett JP, van der Ploeg R, Eijlander RT, Nenninger A, Mendel S, Rozeboom R, et al: The Twin-Arginine Translocation (Tat) systems from *Bacillus subtilis* display a conserved mode of complex organisation and similar substrate recognition requirements. *FEBS J* 2009, 276:232–243.
35. Jongbloed JDH, Antelmann H, Hecker M, Nijland R, Bron S, Airaksinen U, et al: Selective contribution of the twin-arginine translocation pathway to protein secretion in *Bacillus subtilis*. *J Biol Chem* 2002, 277:44068–44078.
36. Smith H, de Jong A, Bron S, Venema G: Characterization of signal-peptide-coding regions selected from the *Bacillus subtilis* chromosome. *Gene* 1988, 70:351–61.
37. Caspers M, Brockmeier U, Degering C, Eggert T, Freudl R: Improvement of Sec-dependent secretion of a heterologous model protein in *Bacillus subtilis* by saturation mutagenesis of the N-domain of the AmyE signal peptide. *Appl Microbiol Biotechnol* 2010, 86:1877–1885.
38. Goosens VJ, Otto A, Glasner C, Monteferante CG, Van der Ploeg R, Hecker M: Novel twin-arginine translocation pathway-dependent phenotypes of *Bacillus subtilis* unveiled by quantitative proteomics. *J Proteome Res*, . in press.
39. Zweers JC, Barák I, Becher D, Driessens AJM, Hecker M, Kontinen VP, et al: Towards the development of *Bacillus subtilis* as a cell factory for membrane proteins and protein complexes. *Microb Cell Fact* 2008, 7:10.
40. Kakeshita H, Kageyama Y, Ara K, Ozaki K, Nakamura K: Enhanced extracellular production of heterologous proteins in *Bacillus subtilis* by deleting the C-terminal region of the SecA secretory machinery. *Mol Biotechnol* 2010, 46:250–257.

41. Diao L, Dong Q, Xu Z, Yang S, Zhou J, Freudl R: Functional implementation of the posttranslational SecB-SecA protein-targeting pathway in *Bacillus subtilis*. *Appl Environ Microbiol* 2012, 78:651–659.
42. Kouwen TRHM, van Dijl JM: Applications of thiol-disulfide oxidoreductases for optimized in vivo production of functionally active proteins in *Bacillus*. *Appl Microbiol Biotechnol* 2009, 85:45–52.
43. Kouwen TRHM, van Dijl JM: Interchangeable modules in bacterial thiol-disulfide exchange pathways. *Trends Microbiol* 2009, 17:6–12.
44. Kouwen TRHM, Dubois JYF, Freudl R, Quax WJ, van Dijl JM: Modulation of thiol-disulfide oxidoreductases for increased production of disulfide bond-containing proteins in *Bacillus*. *Appl Environ Microbiol* 2008, 74:7536–7545.
45. Wu XC, Lee W, Tran L, Wong SL: Engineering a *Bacillus subtilis* expression-secretion system with a strain deficient in six extracellular proteases. *J Bacteriol* 1991, 173:4952–4958.
46. Wu SC, Yeung JC, Duan Y, Ye R, Szarka SJ, Habibi HR, et al: Functional production and characterization of a fibrin-specific single-chain antibody fragment from *Bacillus subtilis*: effects of molecular chaperones and a wall-bound protease on antibody fragment production. *Appl Environ Microbiol* 2002, 68:3261–3269.
47. Westers H, Braun PG, Westers L, Antelmann H, Hecker M, Jongbloed JDH, et al: Genes involved in SkfA killing factor production protect a *Bacillus subtilis* lipase against proteolysis. *Appl Environ Microbiol* 2005, 71:1899–1908.
48. Westers L, Dijkstra DS, Westers H, van Dijl JM, Quax WJ: Secretion of functional human interleukin-3 from *Bacillus subtilis*. *J Biotechnol* 2006, 123:211–224.
49. Antelmann H, Yamamoto H, Sekiguchi J, Hecker M: Stabilization of cell wall proteins in *Bacillus subtilis*: a proteomic approach. *Proteomics* 2002, 2:591–602.
50. Westers L, Westers H, Zanen G, Antelmann H, Hecker M, Noone D, et al: Genetic or chemical protease inhibition causes significant changes in

- the *Bacillus subtilis* exoproteome. *Proteomics* 2008, 8:2704–2713.
51. Yang H, Liu L, Li J, Du G, Chen J: Heterologous expression, biochemical characterization, and overproduction of alkaline α -amylase from *Bacillus alcalophilus* in *Bacillus subtilis*. *Microb Cell Fact* 2011, 10:77.
52. Krishnappa L, Monteferrante CG, van Dijl JM: Degradation of the twinarginine translocation substrate YwbN by extracytoplasmic proteases of *Bacillus subtilis*. *Appl Environ Microbiol* 2012, 78:7801–7804.
53. Goelzer A, Bekkal Brikci F, Martin-Verstraete I, Noirot P, Bessières P, Aymerich S, et al: Reconstruction and analysis of the genetic and metabolic regulatory networks of the central metabolism of *Bacillus subtilis*. *BMC Syst Biol* 2008, 2:20.
54. Goelzer A, Fromion V: Bacterial growth rate reflects a bottleneck in resource allocation. *Biochim Biophys Acta* 2011, 1810:978–88.
55. Goelzer A, Fromion V, Scorletti G: Cell design in bacteria as a convex optimization problem. *Automatica* 2011, 47:1210–1218.
56. Bansal AK: Bioinformatics in microbial biotechnology—a mini review. *Microb Cell Fact* 2005, 4:19.
57. Papagianni M: Recent advances in engineering the central carbon metabolism of industrially important bacteria. *Microb Cell Fact* 2012, 11:50.
58. Li S, Huang D, Li Y, Wen J, Jia X: Rational improvement of the engineered isobutanol-producing *Bacillus subtilis* by elementary mode analysis. *Microb Cell Fact* 2012, 11:101.
59. Westers H, Dorenbos R, van Dijl JM, Kabel J, Flanagan T, Devine KM: Genome engineering reveals large dispensable regions in *Bacillus subtilis*. *Mol Biol Evol* 2003, 20:2076–2090.
60. Morimoto T, Kadoya R, Endo K, Tohata M, Sawada K, Liu S: Enhanced recombinant protein productivity by genome reduction in *Bacillus subtilis*. *DNA Res* 2008, 15:73–81.
61. Manabe K, Kageyama Y, Morimoto T, Ozawa T, Sawada K, Endo K: Combined effect of improved cell yield and increased specific

productivity enhances recombinant enzyme production in genome-reduced *Bacillus subtilis* strain MGB874. *Appl Environ Microbiol* 2011,

77:8370–81.

62. Manabe K, Kageyama Y, Tohata M, Ara K, Ozaki K, Ogasawara N: High external pH enables more efficient secretion of alkaline α -amylase AmyK38 by *Bacillus subtilis*. *Microb Cell Fact* 2012, 11:74.

63. Tanaka K, Henry CS, Zinner JF, Jolivet E, Cohoon MP, Xia F: Building the repertoire of dispensable chromosome regions in *Bacillus subtilis* entails major refinement of cognate large-scale metabolic model. *Nucleic Acids Res* 2013, 41:687–699.

64. le Tam T, Antelmann H, Eymann C, Albrecht D, Bernhardt J, Hecker M: Proteome signatures for stress and starvation in *Bacillus subtilis* as revealed by a 2-D gel image color coding approach. *Proteomics* 2006, 6:4565–85.

65. Kabisch J, Thürmer A, Hübel T, Popper L, Daniel R, Schweder T: Characterization and optimization of *Bacillus subtilis* ATCC 6051 as an expression host. *J Biotechnol* 2012, in press.

66. Marciniaik BC, Trip H, Van-der Veen PJ, Kuipers OP: Comparative transcriptional analysis of *Bacillus subtilis* cells overproducing either secreted proteins, lipoproteins or membrane proteins. *Microb Cell Fact* 2012, 11:66.

67. Toymentseva AA, Schrecke K, Sharipova MR, Mascher T: The LIKE system, a novel protein expression toolbox for *Bacillus subtilis* based on the lal promoter. *Microb Cell Fact* 2012, 11:143